

Computing Special Values of Partial Zeta Functions

Gautam Chinta¹, Paul E. Gunnells¹, and Robert Sczech²

¹ Dept. of Mathematics, Columbia University
New York, NY 10027, USA

² Dept. of Mathematics and Computer Science, Rutgers University
Newark, NJ 07102–1811, USA

Abstract. We discuss computation of the special values of partial zeta functions associated to totally real number fields. The main tool is the *Eisenstein cocycle* Ψ , a group cocycle for $GL_n(\mathbb{Z})$; the special values are computed as periods of Ψ , and are expressed in terms of generalized Dedekind sums. We conclude with some numerical examples for cubic and quartic fields of small discriminant.

1 Introduction

Let K/\mathbb{Q} be a totally real number field of degree n with ring of integers \mathcal{O}_K , and let $U \subset \mathcal{O}_K^\times$ be the subgroup of totally positive units. Let $\mathfrak{f}, \mathfrak{b} \subset \mathcal{O}_K$ be relatively prime ideals. Then the *partial zeta function* associated to these data is defined by

$$\zeta_{\mathfrak{f}}(\mathfrak{b}, s) := \sum_{\mathfrak{a} \sim \mathfrak{b}} N(\mathfrak{a})^{-s},$$

where $\mathfrak{a} \sim \mathfrak{b}$ means $\mathfrak{a}\mathfrak{b}^{-1} = (\alpha)$, where α is a totally positive number in $1 + \mathfrak{f}\mathfrak{b}^{-1}$. According to a classical result of Klingen and Siegel [10], the special values $\zeta_{\mathfrak{f}}(\mathfrak{b}, k)$ are rational for nonpositive integers k . Moreover, the values $\zeta_{\mathfrak{f}}(\mathfrak{b}, 0)$ are especially important because of their connection with the Brumer-Stark conjecture and the Leopoldt conjecture [7,6,3,8,11].

In [9], one of us (RS) gave a cohomological interpretation of these special values by showing that they can be computed in finite terms as periods of the *Eisenstein cocycle*. This is a cocycle $\Psi \in H^{n-1}(GL_n(\mathbb{Z}); \mathcal{M})$, where \mathcal{M} is a certain $GL_n(\mathbb{Z})$ -module. Then two of us (PEG and RS) showed in [5] that the Eisenstein cocycle is an effectively computable object. More precisely, using the cocycle one can express $\zeta_{\mathfrak{f}}(\mathfrak{b}, k)$ as a finite sum of *generalized Dedekind sums*, and the latter can be effectively computed by a continued-fraction algorithm that uses a generalization of the classical Dedekind-Rademacher reciprocity law.

In this note we describe an ongoing project to build a database of $\zeta_{\mathfrak{f}}(\mathfrak{b}, 0)$ for various fields K and ideals $\mathfrak{f}, \mathfrak{b}$. We recall the definition of the Eisenstein cocycle and its relation to the special values (§2), and discuss the effective computation of Dedekind sums (§3). We conclude with examples of special values for some fields of degree 3 and 4 (§4).

2 Dedekind Sums and the Eisenstein Cocycle

2.1

Let σ be a square matrix with integral columns $\sigma_j \in \mathbb{Z}^n$ ($j = 1, \dots, n$), and let $L \subset \mathbb{Z}^n$ be a lattice of rank $r \geq 1$. Let $v \in \mathbb{Q}^n$, and let $e \in \mathbb{Z}^n$ with $e_j \geq 1$. Then the *Dedekind sum* S associated to the data (L, σ, e, v) is defined by

$$S = S(L, \sigma, e, v) := \sum'_{x \in L} \mathbf{e}(\langle x, v \rangle) \frac{\det \sigma}{\langle x, \sigma_1 \rangle^{e_1} \cdots \langle x, \sigma_n \rangle^{e_n}}. \tag{1}$$

Here $\langle x, y \rangle := \sum x_i y_i$ is the usual scalar product on \mathbb{R}^n , $\mathbf{e}(t)$ is the character $\exp(2\pi i t)$, and the prime next to the summation means to omit terms for which the denominator vanishes. The series (1) converges absolutely if all $e_j > 1$, but may only converge conditionally if $e_j = 1$ for some j . In this latter case we can define the sum by the *Q-limit*

$$\sum'_{x \in L} a(x) \Big|_Q := \lim_{t \rightarrow \infty} \left(\sum'_{|Q(x)| < t} a(x) \right), \tag{2}$$

where Q is any finite product of real-valued linear forms on \mathbb{R}^n that doesn't vanish on $\mathbb{Q}^n \setminus \{0\}$. One can precisely determine how the value of (1) depends on Q ([9, Thm. 7]). The sum S is always a rational number times a power of $2\pi i$.

2.2

We recall now the definition of the Eisenstein cocycle Ψ and its relationship with the special values $\zeta_f(\mathbf{b}, k)$. For simplicity, we describe only material necessary to compute the special value at $k = 0$, and refer to [9, 5] for other k .

Let $\mathcal{A} = (A_1, \dots, A_n) \in (GL_n(\mathbb{R}))^n$ be an n -tuple of matrices. For an n -tuple $d = (d_1, \dots, d_n)$ of integers $1 \leq d_i \leq n$, let $\mathcal{A}(d) \subseteq \mathbb{R}^n$ be the subspace generated by all columns A_{ij} such that $j < d_i$. (Here A_{ij} denotes the j th column of the matrix A_i .) Writing $\mathcal{A}(d)^\perp$ for the orthogonal complement of $\mathcal{A}(d)$ in \mathbb{R}^n , we let

$$X(d) = \mathcal{A}(d)^\perp \setminus \bigcup_{i=1}^n \sigma_i^\perp, \quad \text{where } \sigma_i = A_{id_i}. \tag{3}$$

The n -tuple \mathcal{A} determines a decomposition of $\mathbb{R}^n \setminus \{0\}$ into linear strata

$$\bigsqcup_{d \in D} X(d), \tag{4}$$

indexed by the finite set

$$D = D(\mathcal{A}) = \{d \mid X(d) \neq \emptyset\}.$$

Associated to this decomposition is a collection of rational functions $\psi(\mathcal{A})$ on $\mathbb{R}^n \setminus \{0\}$, defined by

$$\psi(\mathcal{A})(x) = \frac{\det(\sigma_1, \dots, \sigma_n)}{\langle x, \sigma_1 \rangle \cdots \langle x, \sigma_n \rangle}, \quad \text{if } x \in X(d).$$

Note that $\psi(\mathcal{A})(x)$ is well-defined by the construction of $X(d)$.

Let $v \in \mathbb{R}^n$, and let Q be defined as in §2.1. Then the *Eisenstein cocycle* Ψ is defined as

$$\Psi = \Psi(\mathcal{A})(Q, v) := (2\pi i)^{-n} \sum_{x \in \mathbb{Z}^n} e(\langle x, v \rangle) \psi(\mathcal{A})(x) \Big|_Q.$$

One can show that Ψ is a homogeneous $(n - 1)$ -cocycle for $GL_n(\mathbb{Z})$. Furthermore, we can express Ψ in terms of Dedekind sums

$$\Psi(\mathcal{A})(Q, v) = (2\pi i)^{-n} \sum_{d \in D} S(L(d), \sigma, \mathbf{1}, v) \Big|_Q, \tag{5}$$

where σ is the matrix with columns A_{id_i} , ($i = 1, \dots, n$), $L(d)$ is the lattice $\mathcal{A}(d)^\perp \cap \mathbb{Z}^n$, and $\mathbf{1}$ is the vector $(1, \dots, 1)$.

2.3

Now we describe how Ψ can be used to compute special values. Let W be a \mathbb{Z} -basis for the fractional ideal $\mathfrak{fb}^{-1} = \sum \mathbb{Z}W_j$, and let W^* be the dual basis with respect to the trace form. Via the n real embeddings τ_i , $i = 1, \dots, n$, any $x \in K$ determines a row vector $(\tau_1(x), \dots, \tau_n(x))$. Hence we may identify W with a matrix in $GL_n(\mathbb{R})$: the j th row of this matrix is the image of the j th basis element of W . Let

$$Q(X) = \prod_i \sum_j X_j(\tau_i(W_j^*)),$$

and let $v \in \mathbb{Q}^n$ be defined by $v_j = \text{Tr}(W_j^*)$.

Let $\nu = n - 1$, and let $\varepsilon_1, \dots, \varepsilon_\nu$ be a basis for the totally positive units U . Using the regular representation ρ with respect to the basis W , we identify the units ε_j with elements $A_j = \rho(\varepsilon_j)^t \in GL_n(\mathbb{Z})$. Using the bar notation

$$[A_1 | \cdots | A_\nu] := (1, A_1, A_1 A_2, \dots, A_1 \cdots A_\nu) \in (GL_n(\mathbb{Z}))^\nu,$$

we have the following proposition expressing the zeta values in terms of the Eisenstein cocycle:

Proposition 1. [9, 5] *Let $U_{\mathfrak{f}}$ be the subgroup $U \cap (1 + \mathfrak{f})$, and let π run through all permutations of $\{1, \dots, \nu\}$. Then*

$$\zeta_{\mathfrak{f}}(\mathfrak{b}, 0) = \eta \sum_{\varepsilon \in U/U_{\mathfrak{f}}} \sum_{\pi} \text{sgn}(\pi) \Psi([A_{\pi(1)} | \cdots | A_{\pi(\nu)}])(Q, \rho(\varepsilon)^t v).$$

Here $\eta = \pm 1$ is defined by

$$\eta = (-1)^\nu \text{sgn}(\det W) \text{sgn}(R),$$

where $R = \det(\log \tau_j(\varepsilon_i))$, $1 \leq i, j \leq \nu$.

3 Diagonality and Unimodularity

3.1

We define the *rank* of $S = S(L, \sigma, e, v)$ to be the rank of the lattice L . It is easy to see that after a $GL_n(\mathbb{Q})$ transformation, we may assume that L is the sublattice $Z^\ell \subset \mathbb{Z}^n$ spanned by the first ℓ standard basis vectors, where ℓ is the rank of L . Furthermore, by multiplying by an appropriate rational factor, permuting columns and repeating columns if necessary, we may assume the pair (Z^ℓ, σ) satisfies the following conditions:

- (i) For each column σ_j , the vector of the first ℓ components of σ_j is primitive and integral.
- (ii) If two columns of σ induce proportional linear forms on Z^ℓ , then these two linear forms coincide on Z^ℓ , and are adjacent columns of σ .
- (iii) The vector $e = \mathbf{1}$.

Let $S(Z^\ell, \sigma, \mathbf{1}, v)$ be a Dedekind sum satisfying the three conditions above. Let $\pi: \mathbb{R}^N \rightarrow \mathbb{R}^\ell$ be the projection on the first ℓ components, and let $\pi(\sigma)$ be the $\ell \times n$ matrix with columns $\pi(\sigma_i)$.

Definition 1. *Let $M(\sigma)$ be the set of maximal minors of $\pi(\sigma)$. Then the index of S , denoted $\|S\|$, is defined to be*

$$\max_{\tau \in M(\sigma)} |\det \tau|.$$

A Dedekind sum is unimodular if $\|S\| = 1$.

3.2

Now define a partition

$$\llbracket n \rrbracket = \bigsqcup_{k=1}^s I_k, \quad \ell \leq s \leq n \tag{6}$$

as follows. Put

$$i, j \in I_k \quad \text{if and only if} \quad \pi(\sigma_i) = \pi(\sigma_j).$$

In other words, two elements of $\llbracket n \rrbracket$ are in the same set of the partition if the corresponding columns of σ induce the same linear form on Z^ℓ .

Let $p_k = \#I_k$.

Definition 2. *The vector $p(S) = (p_1, \dots, p_s)$ is called the type of S . A Dedekind sum is called diagonal if $p(S)$ has length ℓ .*

3.3

The virtue of diagonality is that a diagonal Dedekind sum S may be evaluated as a finite sum of products of generalized Bernoulli polynomials. Furthermore, the number of terms in this finite sum is the index of S . Hence diagonal and unimodular Dedekind sums can be evaluated very rapidly.

In general, the Dedekind sums in (5) aren't diagonal. However, we have the following theorem, which is the main result of [5]:

Theorem 1. [5] *Every Dedekind sum $S(L, \sigma, e, v)$ can be expressed as a finite rational linear combination of unimodular diagonal sums. If n , Rank L , and e are fixed, then this expression can be computed in time polynomial in $\log \|S\|$. Moreover, the number of terms in this expression is bounded by a polynomial in $\log \|S\|$.*

The key ingredient in the proof of Theorem 1 is a “reciprocity law” for higher-dimensional Dedekind sums. For any nonzero point $v \in \mathbb{R}^n$, let v^\perp be the hyperplane $\{x \mid \langle v, x \rangle = 0\}$. Let Q be a finite product of real-valued linear forms on \mathbb{R}^n that do not vanish on $\mathbb{Q}^n \setminus \{0\}$.

Proposition 2. *Let $\sigma_0, \dots, \sigma_n \in \mathbb{Z}^n$ be nonzero. For $j = 0, \dots, n$, let σ^j be the matrix with columns $\sigma_0, \dots, \hat{\sigma}_j, \dots, \sigma_n$. Fix a lattice $L \subseteq \mathbb{Z}^n$, and assume $e = \mathbf{1}$. Then for any $v \in \mathbb{R}^n$, we have the following identity among Dedekind sums:*

$$\sum_{j=0}^n (-1)^j S(L, \sigma^j, \mathbf{1}, v) \Big|_Q = \sum_{j=0}^n (-1)^j S(L \cap \sigma_j^\perp, \sigma^j, \mathbf{1}, v) \Big|_Q. \tag{7}$$

We refer to [5] for proofs of the above statements. Here, in the following two sections, we show how Theorem 1 is applied with a rank 2 example. For simplicity we ignore issues of convergence, and merely remark that all of our manipulations with sums are compatible with the Q -limit process (2).

3.4

Let $L \subset \mathbb{Z}^4$ be the lattice spanned by the first two standard basis vectors, and let

$$\sigma = \begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 2 & 2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad e = \mathbf{1}, \quad \text{and } v = (0, 0, 0, 0).$$

Hence $S(L, \sigma, e, v)$ denotes the absolutely convergent sum

$$\sum'_{(x,y) \in \mathbb{Z}^2} \frac{1}{xy(x+2y)^2},$$

where the prime on the summation indicates that we omit the terms (x, y) for which x, y or $x + 2y$ vanish. This sum isn't diagonal, since σ induces 3 different

linear forms on L instead of 2. Note also that the last two rows of σ don't affect the value of the sum; this observation will play an essential role when we apply Proposition 2 to simplify S .

To diagonalize S , we begin with the identity of rational functions

$$\frac{1}{xy(x+2y)^2} = \frac{1}{y(x+2y)^3} + \frac{2}{x(x+2y)^3}. \tag{8}$$

This is true provided none of the denominators vanishes. The numerators of the functions on the right come from the following identity of linear forms on L :

$$\langle w, (1, 2, *, *)^t \rangle = \langle w, 1 \cdot (1, 0, *, *)^t + 2 \cdot (0, 1, *, *)^t \rangle, \quad \text{for all } w \in L. \tag{9}$$

Here the stars denote entries that we don't care about, since they don't affect the value of the linear form on L . We want to sum both sides of (8) over pairs $(x, y) \in \mathbb{Z}^2$ to obtain an identity among Dedekind sums of the form

$$\sum'_{(x,y) \in \mathbb{Z}^2} \frac{1}{xy(x+2y)^2} = \sum'_{(x,y) \in \mathbb{Z}^2} \frac{1}{y(x+2y)^3} + \sum'_{(x,y) \in \mathbb{Z}^2} \frac{2}{x(x+2y)^3}. \tag{10}$$

However, as written (10) is incorrect. The identity (8) only holds if none of x , y , or $x+2y$ vanish, but the sums on the right of (10) include some of these terms (for instance, the first sum on the right of (10) contains terms (x, y) with $x=0$). We account for this by subtracting two rank 1 Dedekind sums from the right of (10) as ‘‘correction terms’’:

$$\begin{aligned} \sum'_{(x,y) \in \mathbb{Z}^2} \frac{1}{xy(x+2y)^2} &= \sum'_{(x,y) \in \mathbb{Z}^2} \frac{1}{y(x+2y)^3} + \sum'_{(x,y) \in \mathbb{Z}^2} \frac{2}{x(x+2y)^3} \\ &\quad - \sum'_{\substack{(x,y) \in \mathbb{Z}^2 \\ x=0}} \frac{1}{y(x+2y)^3} - \sum'_{\substack{(x,y) \in \mathbb{Z}^2 \\ y=0}} \frac{1}{x(x+2y)^3} \\ &= \sum'_{(x,y) \in \mathbb{Z}^2} \frac{1}{y(x+2y)^3} + \sum'_{(x,y) \in \mathbb{Z}^2} \frac{2}{x(x+2y)^3} \\ &\quad - \sum'_{y \in \mathbb{Z}} \frac{1}{8y^4} - \sum'_{x \in \mathbb{Z}} \frac{2}{x^4}. \end{aligned} \tag{11}$$

Note that all of the sums on the right of (11) are now diagonal.

This equation is precisely an instance of the reciprocity law (Proposition 2). To see this, apply the law with $\sigma_0, \dots, \sigma_4$ the columns of the matrix

$$\begin{pmatrix} 1 & 0 & 1 & 1 & 1 \\ 0 & 1 & 2 & 2 & 2 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix},$$

and with L and v as above. Note that $\sigma_4 = \sigma_0 + 2\sigma_1$, which is exactly the linear relation (9). The three rank 2 sums (respectively, the two rank 1 sums) are the

left (resp. right) of (7). All other sums vanish identically, either from the linear dependence among the σ_i , or because all terms are meaningless. Notice how we used that the last two rows of the σ_i have no effect on the sum: this enabled us to introduce a linear dependence among the σ_i that killed some of the nondiagonal sums.

To diagonalize a general Dedekind sum $S(L, \sigma, \mathbf{1}, v)$, one considers the configuration $C \subset \mathbb{R}^n$ of linear subspaces consisting of $(L \otimes \mathbb{R})^\perp$ and the spaces generated by the points $\sigma_1, \dots, \sigma_n$. One shows by investigating the geometry of C that a point σ_0 can be found such that when Proposition 2 is applied with the tuple $(\sigma_0, \dots, \sigma_n)$, the resulting Dedekind sums are “closer” to diagonality in a certain sense. It may take several applications of Proposition 2 to express a Dedekind sum as a linear combination of diagonal sums.

3.5

The second rank 2 sum on the right of (11) has index 2. We will show how to make this sum unimodular. Let $\sigma_0, \dots, \sigma_4$ be the columns of the matrix

$$\begin{pmatrix} 1 & 1 & 1 & 1 & 0 \\ 0 & 2 & 2 & 2 & 1 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{pmatrix},$$

and let L and v be as above. Then an application of Proposition 2 yields

$$\sum'_{(x,y) \in \mathbb{Z}^2} \frac{2}{x(x+2y)^3} = \sum'_{(x,y) \in \mathbb{Z}^2} \frac{-1}{y(x+2y)^3} + \sum'_{(x,y) \in \mathbb{Z}^2} \frac{1}{x(x+2y)^2 y} + \sum'_{y \in \mathbb{Z}} \frac{1}{8y^4} + \sum'_{x \in \mathbb{Z}} \frac{2}{x^4}. \tag{12}$$

Now all the terms on the right of (12) are diagonal and unimodular except for the second rank 2 sum. In fact, this sum is no longer diagonal. However, one further application of Proposition 2 will make this sum diagonal and unimodular. Hence we will have succeeded in expressing the original sum as a finite linear combination of diagonal, unimodular Dedekind sums.

In proceed in general, one must be able to construct the “index-reducing” vector σ_4 as above. An easy argument using Minkowski’s Theorem guarantees the existence of such a vector [1]. To construct this vector in practice, one may use *LLL*-reduction and [4, Conjecture 3.9].

4 Examples

Here we present some numerical examples. For simplicity we compute $\zeta = \zeta_{\mathfrak{f}}(\mathfrak{b}, 0)$, where $\mathfrak{f} = N\mathcal{O}_K$ for various rational integers N , and $\mathfrak{b} = \mathcal{O}_K$. These fields are the first entries in the tables of totally real fields with small discriminant, available from [2].

Cubic fields

- $K = \mathbb{Q}(\theta)$, where $\theta^3 + \theta^2 - 2\theta - 1 = 0$ (discriminant 49).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
		4	1	7	2	10	16	13	-1	16	-9	19	18	22	-8
2	0	5	2	8	3	11	-10	14	-24	17	-26	20	19	23	10
3	2	6	8	9	2	12	5	15	14	18	8	21	6	24	23

- $K = \mathbb{Q}(\theta)$, where $\theta^3 - 3\theta - 1 = 0$ (discriminant 81).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
		4	1	7	14	10	-16	13	2	16	-13	19	11	22	72
2	0	5	-2	8	7	11	-6	14	0	17	13	20	11	23	62
3	2/3	6	8/3	9	2/3	12	11/3	15	14/3	18	-64/3	21	14/3	24	29/3

- $K = \mathbb{Q}(\theta)$, where $\theta^3 + \theta^2 - 3\theta - 1 = 0$ (discriminant 148).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
		4	1	7	2	10	-2	13	-22	16	7	19	82	22	68
2	0	5	-4	8	3	11	-18	14	-20	17	100	20	4	23	12
3	2	6	4	9	-10	12	5	15	42	18	-32	21	-78	24	23

- $K = \mathbb{Q}(\theta)$, where $\theta^3 - \theta^2 - 4\theta - 1 = 0$ (discriminant 169).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
		4	3	7	6	10	8	13	2	16	25	19	-238	22	160
2	0	5	1	8	5	11	-6	14	56	17	6	20	-8	23	386
3	2	6	8	9	26	12	11	15	17	18	52	21	-10	24	89

- $K = \mathbb{Q}(\theta)$, where $\theta^3 - 4\theta - 1 = 0$ (discriminant 229).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
		4	2	7	18	10	-24	13	60	16	19	19	26	22	8
2	0	5	-2	8	7	11	-4	14	24	17	8	20	38	23	202
3	2	6	0	9	-10	12	18	15	56	18	-16	21	96	24	51

- $K = \mathbb{Q}(\theta)$, where $\theta^3 - \theta^2 - 4\theta + 3 = 0$ (discriminant 257).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
		4	6	7	16	10	4	13	10	16	5	19	42	22	72
2	0	5	3	8	9	11	-6	14	0	17	-26	20	12	23	-112
3	2	6	8	9	-4	12	8	15	21	18	-16	21	-12	24	58

Quartic fields

- $K = \mathbb{Q}(\theta)$, where $\theta^4 - \theta^3 - 3\theta^2 + \theta + 1 = 0$ (discriminant 725).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
		4	1	7	-4	10	0	13	-20	16	35	19	-32	22	32
2	0	5	4	8	7	11	-2	14	-16	17	-92	20	0	23	-12
3	4	6	0	9	-44	12	0	15	0	18	-320	21	-30	24	-84

- $K = \mathbb{Q}(\theta)$, where $\theta^4 - \theta^3 - 4\theta^2 + 4\theta + 1 = 0$ (discriminant 1125).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
2	0	3	4/5	5	4/3	7	124/3	9	-116/5	11	72	13	676/3	15	4
		4	2	6	0	8	7	10	128	12	0	14	144	16	51

- $K = \mathbb{Q}(\theta)$, where $\theta^4 - 6\theta^2 + 4 = 0$ (discriminant 1600).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
2	1	4	1/2	7	4	10	4	13	104	16	45/2	19	224	22	136
3	4	5	4	8	5/2	11	-4	14	-16	17	-84	20	-9	23	60
		6	0	9	16	12	24	15	6	18	128	21	222	24	15

- $K = \mathbb{Q}(\theta)$, where $\theta^4 - 4\theta^2 - \theta + 1 = 0$ (discriminant 1957).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
2	0	3	4	5	20	7	-8	9	52	11	4
		4	1	6	0	8	11	10	0	12	3

- $K = \mathbb{Q}(\theta)$, where $\theta^4 - 5\theta^2 + 5 = 0$ (discriminant 2000).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
2	2/5	4	11/5	7	52	10	0	13	-32	16	31/10	19	412/5	22	568
3	4	5	0	8	71/10	11	4	14	16	17	296	20	30		
		6	0	9	4	12	12	15	340	18	-112	21	428		

- $K = \mathbb{Q}(\theta)$, where $\theta^4 - 4\theta^2 + 2 = 0$ (discriminant 2048).

N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$	N	$N \cdot \zeta$
2	1/2	3	4	5	-28	7	8	9	-20
		4	1/4	6	-10	8	9/4	10	19

References

1. A. Ash and L. Rudolph, *The modular symbol and continued fractions in higher dimensions*, *Invent. Math.* **55** (1979), 241–250.
2. J. Buchmann, D. Ford, M. Pohst, M. Olivier, and F. Diaz y Diaz, *Tables of number fields of low degree*, <ftp://megrez.math.u-bordeaux.fr/pub/numberfields/>.
3. D. S. Dummit and D. R. Hayes, *Checking the p-adic Stark conjecture when p is Archimedean*, *Algorithmic number theory* (Talence, 1996), Springer, Berlin, 1996, pp. 91–97.
4. P. E. Gunnells, *Computing Hecke eigenvalues below the cohomological dimension*, *J. Experimental Math.* (to appear), 2000.
5. P. E. Gunnells and R. Schech, *Evaluation of Dedekind sums, Eisenstein cocycles, and special values of L-functions*, preprint, 1999.
6. D. R. Hayes, *Brumer elements over a real quadratic base field*, *Exposition. Math.* **8** (1990), no. 2, 137–184.
7. ———, *The partial zeta functions of a real quadratic number field evaluated at $s = 0$* , *Number theory* (Banff, AB, 1988), de Gruyter, Berlin, 1990, pp. 207–226.

8. ———, *Aligning Brumer-Stark elements into a Hecke character* (working paper, preprint, 1998).
9. R. Sczech, *Eisenstein group cocycles for GL_n and values of L -functions*, *Invent. Math.* **113** (1993), no. 3, 581–616.
10. C. S. Siegel, *Über die Fourierschen Koeffizienten von Modulformen*, *Nachrichten der Akademie der Wissenschaften in Göttingen, Mathematisch-physikalische Klasse* (1970), no. 3, 15–56.
11. L. C. Washington, *Introduction to cyclotomic fields*, second ed., *Graduate Texts in Mathematics*, no. 83, Springer-Verlag, 1997.